

Total No. of Printed Pages—7

**3 SEM TDC MTMH (CBCS) C 5**

**2 0 2 1**

( Held in January/February, 2022 )

**MATHEMATICS**

( Core )

Paper : C-5

( Theory of Real Functions )

Full Marks : 80

Pass Marks : 32

Time : 3 hours

*The figures in the margin indicate full marks  
for the questions*

1. (a) Define limit of function at a point. 1  
(b) Evaluate the following limits (any one) : 2

(i)  $\lim_{x \rightarrow 2} \sqrt{\frac{2x+1}{x+3}}$

(ii)  $\lim_{x \rightarrow 1} \frac{x-1}{\sqrt{x+3}-2}$

( 2 )

- (c) If  $f : A \rightarrow R$  and if  $c$  is a cluster point of  $A$ , then prove that  $f$  can have only one limit at  $c$ . 3

2. (a) Write the type of discontinuity if

$$\lim_{x \rightarrow c^+} f(x) \neq \lim_{x \rightarrow c^-} f(x) \quad 1$$

- (b) When does a function  $f$  continuous on a set? 2

- (c) Investigate for the point of discontinuity : 4

$$f(x) = \begin{cases} 1; & \text{if } x \text{ is rational} \\ 0; & \text{if } x \text{ is irrational} \end{cases}$$

Or

Let  $A, B \subseteq R$  and let  $f : A \rightarrow R$  and  $g : B \rightarrow R$  be functions such that  $f(A) \subseteq B$ . If  $f$  is continuous at a point  $c \in A$  and  $g$  is continuous at  $b = f(c) \in B$ ; then prove that composition  $g \circ f : A \rightarrow R$  is continuous at  $c$ .

( 3 )

- (d) Let  $A \subseteq R$ , let  $f : A \rightarrow R$  and let  $|f|$  be defined by  $|f|(x) = |f(x)|$  for  $x \in A$  and  $f$  is continuous at a point  $c \in A$ . Prove that  $|f|$  is continuous at  $c$ . 3

Or

Discuss the continuity of  $f(x) = |x-1| + |x-2|$  in the interval  $[0, 3]$ .

3. (a) State location of roots theorem. 1

- (b) State and prove intermediate value theorem. 4

- (c) Find the roots of the equation  $x^3 - x - 1 = 0$  between 1 and 2 by using location of roots (bisection method) theorem. 4

Or

Let  $I$  be a closed bounded interval and let  $f : I \rightarrow R$  be continuous on  $I$ , then prove that the set  $f(I) = \{f(x) : x \in I\}$  is a closed bounded interval.

4. (a) Write the non-uniformity continuity criteria (any one). 1

( 4 )

- (b) Show that a function  $f : R \rightarrow R$  given by  $f(x) = x^2$  is not uniformly continuous on  $R$ . 4

Or

If  $f$  and  $g$  are each uniformly continuous on  $R$ , then prove that composite function  $f \circ g$  is also uniformly continuous on  $R$ .

5. (a) Find :  $\frac{d}{dx}(\tan x^2)$  1
- (b) State Caratheodory's theorem. 2
- (c) If  $f$  is continuous on the closed interval  $I = [a, b]$  and  $f$  is differentiable on the open interval  $(a, b)$  and  $f'(x) = 0$  for all  $x \in (a, b)$ , prove that  $f$  is constant on  $I$ . 3
6. (a) Define relative maximum and relative minimum at a point on an interval. 2
- (b) State and prove Rolle's theorem. 1+3=4

( 5 )

- (c) Apply the mean value theorem to prove the following (any one) : 4

(i)  $e^x \geq 1+x$  for  $x \in R$

(ii)  $\frac{b-a}{1+b^2} < \tan^{-1} b - \tan^{-1} a < \frac{b-a}{1+a^2}$   
for  $a < b$

7. (a) Show that  $f(x) = x^3 - 3x^2 + 3x + 2$  is strictly increasing for every value of  $x \in R$  except 1. 2
- (b) Let  $I \subseteq R$  be an interval, let  $f : I \rightarrow R$ , let  $c \in I$  and assume that  $f$  has a derivative at  $c$  and  $f'(c) > 0$ , then there is a number  $\delta > 0$ . Prove that  $f(x) > f(c)$  for  $x \in I$  and  $c < x < c + \delta$ . 3
- (c) Examine the validity of mean value theorem for the function  $f(x) = 2x^2 - 7x + 10$  on  $[2, 5]$ . 4

Or

If  $f$  is differentiable on  $I = [a, b]$  and if  $k$  is a number between  $f'(a)$  and  $f'(b)$ , then prove that there exists at least one point  $c$  in  $(a, b)$ , where  $f'(c) = k$ .

8. (a) Write the remainder after  $n$  terms of Taylor's theorem in Lagrange's form. 1
- (b) Write the statement of Cauchy's mean value theorem. 2
- (c) Deduce from Cauchy's mean value theorem  $f(b) - f(a) = \xi f'(\xi) \log \frac{b}{a}$ , where  $f(x)$  is continuous and differentiable in  $[a, b]$  and  $a < \xi < b$ . 3
- (d) State and prove Taylor's theorem with Cauchy's form of remainder. 6

Or

Find the approximate value of  $\sqrt[3]{1+x}$ ,  $x > -1$  by using Taylor's theorem with  $n = 2$ .

9. (a) Write the necessary condition for a function  $f(x)$  to have relative extremum at  $x = c$ . 1
- (b) Determine whether or not  $x = 0$  is a point of relative extremum of  $f(x) = \sin x - x$ . 2
- (c) Define convex function. 2

- (d) Using Maclaurin's series, expand the following in an infinite series in powers of  $x$  (any two) :  $4 \times 2 = 8$

(i)  $\log(1+x)$

(ii)  $\cos x$

(iii)  $\frac{1}{ax+b}$

\*\*\*